

Review Article

Human factors in anaesthesia: a narrative review

F. E. Kelly,¹  C. Frerk,^{2,3} C. R. Bailey,⁴ T. M. Cook,^{1,5} K. Ferguson,⁶ R. Flin,⁷ K. Fong,^{8,9} P. Groom,¹⁰ C. John,¹¹ A. R. Lang,¹² T. Meek,¹³ K. L. Miller,¹⁴ L. Richmond,¹⁵ N. Sevdalis¹⁶ and M. R. Stacey¹⁷

1 Consultant, Department of Anaesthesia and Intensive Care Medicine, Royal United Hospitals Bath NHS Foundation Trust, Bath, UK

2 Consultant, Department of Anaesthesia and Critical Care, Northampton General Hospital, Northampton, UK, and Difficult Airway Society Professor of Anaesthesia and Airway Management

3 Honorary Visiting Professor, University of Leicester, College of Life Sciences/Leicester Medical School, Leicester, UK

4 Consultant, Department of Anaesthesia, Guy's and St Thomas' NHS Foundation Trust, London, UK

5 Honorary Professor, School of Medicine, Bristol University, Bristol, UK

6 Consultant, Department of Anaesthesia, Aberdeen Royal Infirmary, Aberdeen, UK, and Past President, Association of Anaesthetists

7 Professor, School of Psychology, Aberdeen Business School, Robert Gordon University, Aberdeen, UK

8 Consultant, Department of Anaesthesia, University College London Hospitals NHS Foundation Trust, London, UK

9 Professor, Department of Science, Technology, Engineering and Public Policy, University College London, UK

10 Consultant, Department of Anaesthesia, Liverpool University Hospitals NHS Foundation Trust, Liverpool, UK

11 Operating Department Practitioner and Head of Clinical Education, Department of Anaesthesia, University College Hospital's NHS Foundation Trust, London, UK

12 Assistant Professor, Human Factors Research Group, Faculty of Engineering, University of Nottingham, UK, and previous Chair of Healthcare Sector Group, Chartered Institute of Ergonomics and Human Factors

13 Consultant, Department of Anaesthesia, James Cook University Hospital, Middlesbrough, UK, and Vice-president, Association of Anaesthetists

14 Consultant, Department of Anaesthesia, Birmingham Women's and Children's Hospital NHS Foundation Trust, Birmingham, UK and Trainee Committee, Association of Anaesthetists

15 Consultant, Department of Anaesthesia, Swansea Bay University Health Board, Swansea, UK, and previous trainee representative Difficult Airway Society

16 Professor, Centre for Implementation Science, King's College London, UK

17 Consultant, Department of Anaesthesia, Intensive Care and Pain Medicine, University Hospital of Wales, Cardiff, UK, Deputy Dean, Health Education Improvement Wales, and Board Member, Association of Anaesthetists

Summary

Healthcare relies on high levels of human performance, as described by the 'human as the hero' concept. However, human performance varies and is recognised to fall in high-pressure situations, meaning that it is not a reliable method of ensuring safety. Other safety-critical industries embed human factors principles into all aspects of their organisations to improve safety and reduce reliance on exceptional human performance; there is potential to do the same in anaesthesia. Human factors is a broad-based scientific discipline which aims to make it as easy as possible for workers to do things correctly. The human factors strategies most likely to be effective are those which 'design out' the chance of an error or adverse event occurring. When errors or adverse events do happen, barriers are in place to trap them and reduce the risk of progression to patient and/or worker harm. If errors or adverse events are not trapped by these barriers, mitigations are in place to minimise the consequences. Non-technical skills form an important part of human factors barriers and mitigation strategies and include: situation awareness; decision-making; task management; and team working. Human factors principles are not a substitute for proper investment and appropriate staffing levels. Although applying human factors science has the potential to save money in the long term, its proper implementation may require investment before reward can be reaped. This narrative review describes what is known about human factors in anaesthesia to date.

Correspondence to: F. E. Kelly
Email: fiona.kelly@doctors.org.uk

Accepted: 8 November 2022

Keywords: anaesthesia; ergonomics; human factors; non-technical skills education and training; patient safety

This article accompanies guidelines by Kelly et al., *Anaesthesia* 2023; <https://doi.org/10.1111/anae.15941>

Twitter: @fionafionakel; @chrisfrerk; @DocCRBailey; @doctimcook; @KFAberdeen; @rhonaflin; @Kevin_Fong; @clintonejohn; @lanalex83; @drtimmeek; @nemolysis; @NickSevdalis; @airwayman1

Introduction

Humans have physical and cognitive limitations and are fallible. The safety of patients should, therefore, not rely on high levels of individual and team performance but should instead depend on well-designed systems that have been put in place to support staff. Human factors science is part of a scientific principle that accepts and understands that humans are fallible [1, 2], and draws on engineering, psychology, social and physical anthropology to design systems that make it easy for people 'to do the right thing', and difficult or impossible 'to do the wrong thing' [3, 4]. Human factors science is embedded within many safety-critical industries and provides an opportunity to improve safety in anaesthesia and in healthcare in general [2, 4]. The International Ergonomics Association defines human factors as "the scientific discipline concerned with the understanding of interactions among humans and other elements of a system", and "the profession that applies theory, principles, data and methods to design in order to optimise human wellbeing and overall system performance" [1].

Human factors strategies can be grouped together into four domains and ordered in a hierarchy of controls model according to their likely effectiveness (Fig. 1) [5]. Providing well-designed work environments and equipment to

decrease the chance of errors occurring is the strategy most likely to be effective and forms the stable base of the pyramid. Barriers (physical and administrative) aim to 'trap' potential errors, reducing the chance of them progressing to patient and/or staff harm. Not all barriers are equal in their efficacy and this needs to be understood and considered in their use. Mitigations comprise behaviours and working practices that form rescue systems: these reduce harm to patients and staff if a significant error progresses past the design and barriers described above. Finally, education and training form the top of the pyramid and, while essential for safety, cannot compensate for poor environment, equipment and systems. Regardless of how well-trained staff are, errors are likely to occur if they are asked to work in poorly designed environments with poor working systems in place. It is important that the base of the pyramid is the largest part, with commensurate economic investment to provide a stable and effective foundation for safety [6–8]. The hierarchy of controls model described in this paper is based on the 'error troika' model [9] and general human factors principles, which have been used in healthcare [5]. In comparison, the National Institute for Occupational Safety and Health (NIOSH) hierarchy of controls model [10], developed with the primary aim of protecting staff from hazards rather than optimising work overall, is not as transferable a model to healthcare [11].

A well-known patient safety concept, the 'Swiss cheese model' [12] (Fig. 2), describes a potential hazard travelling through a system, with slices of cheese representing barriers that prevent hazard progression to an adverse event. The holes in the cheese represent unsafe systems and active errors, and only when these holes are aligned can the hazard pass through. With poorly designed healthcare systems, the holes are large, barriers are less effective and too much reliance is placed on individual practitioners consistently performing at very high levels [12, 13]. This illustrates the 'human as the hero' model [12], where exceptional performance is required to maintain safety every day. As safer systems are designed and implemented,




Figure 1 Hierarchy of controls model describing human factors strategies arranged according to their likely effectiveness.




Figure 2 Swiss Cheese model (reproduced with permission from [13]).

the holes become smaller, chance of alignment falls and the risk of the hazard causing an adverse event is decreased.

Impact of sentinel cases on human factors in anaesthesia and wider healthcare

Non-technical skills are an important component of human factors. They include cognitive, social and personal skills that complement technical skills [14, 15] and feature in the barriers and mitigation strategies which reduce the harm and consequences of an adverse event. The value of non-technical skills in anaesthesia has been recognised since the 1990 s [16–18], with the Anaesthetic Non-Technical Skills (ANTS) framework being described in 2003 [18] and the benefits for everyday working and crisis management outlined [17, 18]. Non-technical skill training became much more common following the death of Elaine Bromiley in 2005 [3, 19], with widespread human factors publicity afforded to this by her husband Martin and the anaesthetic profession's response to his work [3, 20]. Martin Bromiley's impact was the dawning of non-technical skills as an issue in anaesthesia safety. This may have led in part to the belief that human factors science is the same as non-technical skills. Bromiley, an airline pilot, emphasised the role of non-technical skills training in improving teamwork, safety and crisis management – such training had been standard in the airline industry since the 1980 s [3, 20]. In addition, he highlighted the importance of a flattened team hierarchy, use of checklists, recognition of situation awareness and strategies to improve it and promoted the integration of human factors training into medical education and healthcare management [3, 20, 21]. Bromiley still regards non-technical skills as critical to safety but recognises that human factors principles around design, systems and work patterns are already embedded at all levels of his industry, providing a firm foundation for the application of good non-technical skills during everyday work [4].

The death of Glenda Logsdail following unrecognised oesophageal intubation in 2020 has raised further

awareness of the role of human factors in patient safety [22]. Glenda's husband, Richard, works in the offshore oil industry and is keen that her story is shared, lessons are identified and that the design of safe systems is prioritised to prevent future deaths [23, 24].

Strategies for improving human factors in anaesthesia

Design of medical equipment

Medical equipment and devices range from simple items (e.g. disposable gloves, hypodermic needles and facemasks) to far more complex devices (e.g. ultrasound machines and anaesthesia workstations).

Well-designed medical equipment can reduce the chance of errors occurring with advantages for patients and staff [25]. One example is videolaryngoscopy which has been shown to improve the visualisation of the glottis [26, 27], improve tracheal intubation success rate [26, 27], and reduce the incidence of failed tracheal intubation [26, 27] and oesophageal intubation [26]. Videolaryngoscopy also reduces mechanical stress on the neck, back and shoulders of the clinician, decreasing the likelihood of worker injury [28] and can prevent the transmission of infections during tracheal intubation by increasing the distance between the patient's and operator's airways [29]. The benefits seen in increasing success rates for tracheal intubation with videolaryngoscopy are particularly noticeable when clinicians are wearing personal protective equipment (PPE) [30].

Medical device and equipment manufacturers are required to perform human factors assessments on their products before releasing them to market, according to international standards [31, 32] and Medicines and Healthcare products Regulatory Agency (MHRA) guidance [33]. At present, manufacturers must demonstrate that such assessments occur during the design process, although this often occurs too late for them to make changes if any unanticipated design flaws are detected. In this situation, manufacturers merely have to demonstrate that they have taken steps to mitigate any associated risk, which could be

as simple as documenting this in the product instruction manual. Performing human factors usability assessments at an earlier stage of product design and development would allow equipment to be redesigned before being marketed. The human factors assessments performed during the design and development stages, and the resulting mitigations, remain the intellectual property of the manufacturers and are not routinely available to individual hospitals. In-house evaluations cannot expect to uncover hazards and difficulties in use that would be revealed by a comprehensive 'use assessment' by a human factors professional, which can result in equipment with potential hazards finding its way into clinical practice [34].

Design of drug ampoules and packaging

All pharmaceutical companies follow international [35, 36], European and MHRA [37] regulations regarding the colour and design of labelling and packaging; however, this does not always provide optimum readability nor ease of use for clinicians, and drug administration errors continue to occur [38–42]. Different drugs produced by the same manufacturer can look strikingly similar (Fig. 3). Conversely, the same drug produced by different manufacturers can have very different ampoule and packaging design (Fig. 3), resulting in the appearance of drugs changing in the event of a supply chain shortage or change in pharmacy purchasing contracts. Improvements in design have the potential to reduce selection errors associated with drugs that look similar. Potential strategies to improve drug ampoule and packaging design are being investigated.

Design and layout of operating theatres

Operating theatre design and layout should make it as easy as possible for the whole team to work effectively while

maintaining patient safety [43]. However, anaesthetists sometimes find themselves working in confined spaces surrounded by cables and pipelines, with the patient monitor difficult to see and infusion devices and urinary catheters awkward to access. There are two principal reasons for this. First relates to operating theatre design. Decisions regarding positioning of fixed features including operating theatre lights, laminar flow hoods, ceiling pendants, pipeline gases, scavenging, suction and power sockets are made when the operating theatre is built. Although this is done in accordance with building regulations, it does not usually consider the way in which the staff will use the space. As a result, staff are forced to adjust their pattern of work to accommodate the operating theatre design. Second is workspace layout. This is the way in which the theatre team arrange moveable equipment within their work area on a daily basis [44]. Compromises are needed between the surgical, anaesthetic and scrub teams who compete for space, a problem that is increasing as the amount and size of operating theatre equipment grows. Human factors tools can be used to identify optimum workspace layouts, and in doing so improve efficiency and reduce the risk of musculoskeletal injuries for staff [43, 44].

Design of equipment storage areas

Well-designed equipment storage areas can reduce the time taken to access equipment in an emergency [45], with examples being standardised layout of cardiac arrest [46] and difficult airway trolleys [47].

Design of safe working practices

Good design of working practices is also essential for ensuring safety. This includes ensuring adequate staffing levels and skill mix, with plans which allow for surges in clinical demand [4]. In addition, staff supervision



Figure 3 Drug ampoules that look similar, but which contain different drugs.

arrangements, lines of accountability, systems for identification of patients who are sick and/or deteriorating, and systems to allow staff to escalate concerns should be in place [4]. An understanding of demand, capacity and 'work as done' by healthcare staff is needed to design systems and working practices that reflect real life as closely as possible [2, 4].

Barriers

Barriers prevent errors that have occurred from progressing to cause harm to patients and staff; they should be incorporated into working practices when there is no reliable method for 'designing out' the potential hazard.

Administrative barriers

Well-designed policies, protocols and standard operating procedures are easier for staff to use and are more likely to be followed accurately with fewer deviations and omitted steps [48–50]. Nationally agreed policies and protocols with standardised content, formatting and language, can help workers moving between different hospitals, reduce variation in practice and may raise standards [51].

Staff who need to rush while working and/or are anxious due to potentially unachievable workloads are more likely to make mistakes [52–57]. Theatre list planning that acknowledges the additional time required for complex patients, difficult procedures and a high turnover of cases reduces time pressures on staff and allows them to plan and prepare for such cases [58]. Making individual and team workloads more manageable results in improved patient safety as well as reducing the risk of cancellations.

Physical barriers

Physical barriers can prevent harm to staff and patients, with examples including protective goggles and locked theatre doors during laser procedures and simple PPE such as gowns, gloves and masks.

Cognitive aids

Cognitive aids are tools designed to guide staff while they are performing a task or group of tasks [59]. The use of cognitive aids can enhance worker performance, including speed, accuracy and fluidity of task management [59, 60]. Examples of anaesthetic cognitive aids include the Association of Anaesthetists quick reference handbook [61] and the Vortex approach for managing an airway emergency [62].

Algorithms are cognitive aids which comprise an ordered sequence of steps with each step depending on the outcome of the previous one. The 2015 DAS guidelines

include examples of well-designed algorithms, with prompts for the team to 'call for help', 'declare a failed intubation', 'stop and think' and 'invite ideas from the team' [47].

Checklists are also cognitive aids, some of which have been shown to reduce postoperative complications and improve surgical outcomes [63–65]. Examples in anaesthesia include the World Health Organization surgical safety checklist [63] and rapid sequence induction checklists [66, 67]. The success of checklists relies on staff being able to use them consistently in the way in which they were designed [64, 65].

Practising non-technical skills during every day working

Non-technical skills are included in the UK Royal College of Anaesthetists' training curriculum [68] and form part of the introduction to the science of human factors for many anaesthetists. Non-technical skills in anaesthesia are well-described by the ANTS system (Fig. 4) [14, 15], and include: communication; leadership and followership [69]; delegation and prioritisation; coping with distractions and fatigue; and managing stress [14]. These can be learned and practised just like technical skills [14, 70].

Situation awareness can be defined as a cognitive skill that involves 'selection and comprehension of information from the world around us to make sense of our work environment'. It has three components: gathering information; comprehension; and anticipation [14]. In plain English, this means knowing what is going on around you, understanding past events, and anticipating what might happen next [14, 72]. Loss of situation awareness is often associated with the limited capacity of human working memory and the brain being susceptible to 'cognitive overload' in high-pressure situations [72, 73]. Factors that reduce situation awareness include: stress; fatigue; lack of expertise; high workload; and multiple distractions. Education and training, including using simulation, has been shown to improve understanding of situation awareness [14] and provides opportunities to practise strategies to maintain it [14].

Decision-making involves identifying and defining a problem, considering options, selecting and implementing one of those options, and then reviewing the outcome [14]. Decision-making can be learned and improved with practise [14] and is impaired by the same factors that reduce situation awareness.

Task management describes the skills required for organising resources and activities to complete a procedure or plan of action. It can be subdivided into planning and preparing, prioritising, maintaining standards and identifying and utilising resources [14, 15, 69].




Figure 4 Anaesthetists Non-Technical Skills (ANTS) system (reproduced with permission from [71]).

Effective teamworking is essential for the practice of safe, high-quality anaesthesia [14, 74]. Team working issues including poorly defined roles, lack of explicit coordination, communication problems and failures to resolve conflict have all been identified as contributing to high profile anaesthetic cases that have not gone to plan [3, 19, 20, 22, 75–77].

Effective communication is a non-technical skill that can be learned, developed and improved [14]. It involves the sender encoding and transmitting the desired information followed by receiver receiving, decoding and understanding the information. ‘Closed loop communication’ or ‘read-back’, where explicit confirmation is given that a message has been received and understood can help to avoid misunderstandings [78, 79].

There are multiple challenges for anaesthetists when developing and using their non-technical skills. Anaesthetists work in high-pressure environments, often caring for patients who are sick in emergency situations. Anaesthetists often need to process large amounts of information in a limited time with a resultant high cognitive load [80]. Multiple interruptions are common [81], with one study reporting an average of 33 distractions per hour in an

operating theatre [82]. Theatre team composition may be different each day and may even change during an operating list. In emergency situations, ‘flash teams’ form [83]: the members of these teams are unlikely to have had a briefing at the start of a shift and may not know each other’s names, roles, skill sets or levels of experience [83]. In all these situations, the use of good non-technical skills can help increase margins of safety.

‘Flattening of the hierarchy’ within a team

Good use of non-technical skills can help to ‘flatten the hierarchy’ within the team, also described as ‘reducing the authority gradient’ [14]. This can help team members to challenge more senior colleagues if they believe that patient safety is at risk [83–85]. A certain degree of hierarchy within a team is necessary for effective leadership and followership. However, a steep hierarchy, and subsequent failure of team members to speak up (or be heard) when they raised concerns, appears to have played a part in high profile deaths in the UK, including Elaine Bromiley [3, 19, 20], Sharon Grierson [76], Peter Saint [77] and Glenda Logsdail [22]. Graded assertiveness tools such as CUSS (“I’m concerned”; “I’m uncomfortable”; “This doesn’t feel safe”;

‘top’ [83, 86] may help empower team members to speak up and provide a structure to use when they do so. These tools work most effectively if senior staff are also trained to ‘listen down’ for the specific phrases used in such tools.

Task fixation and transitioning from one stage of an algorithm to the next

Task fixation describes the situation when an individual or team make multiple unsuccessful attempts at one technique and become ‘stuck’ on one procedure or at one stage of an algorithm [87]. Strategies to encourage staff to transition from one part of an algorithm to the next help reduce the risk of this occurring. The Advanced Life Support algorithm gives time-based transition prompts (recheck pulse every 2 min) as well as outcome-based action prompts (shock or no shock depending on rhythm) to help the resuscitation team move through the correct management steps. Examples of anaesthetic cognitive aids which encourage transitioning can be found in the DAS guidelines, with a specified maximum number of attempts at tracheal intubation and supraglottic airway insertion [47].

Mitigations

Mitigations are systems or actions which reduce the consequences of an error that has occurred, with the aim of preventing or minimising subsequent harm to patients and/or staff. Examples of mitigations familiar to anaesthetists include the ready availability of ‘anaphylaxis boxes’ and the activation of outreach teams in response to early warning scoring systems being triggered. Less obvious examples of mitigations include incident investigation and the associated learning along with strategies and actions to support staff affected following a critical event.

Incident investigation

Incident investigations should ensure that lessons are identified, actions are put in place and their effectiveness is monitored. The investigation techniques used currently in healthcare often scrutinise the actions of staff involved but do not always identify system factors. As a result, investigations may fail to identify robust system changes that could be implemented to prevent the event recurring, and instead just lead to recommendations for additional staff training [88]. A new patient safety incident response framework (PSIRF), based on human factors principles [89], is replacing the existing root cause analysis investigation tools that are currently in use in the UK healthcare system. This framework recommends that critical event investigation teams should include members with human factors training and use human factors tools such as systems engineering

initiative for patient safety (SEIPS) [90]. Another human factors tool validated for use in healthcare investigations is the Yorkshire contributory factors framework [91].

Learning from events that go well as well as ones that do not

Safety-I is defined as a state where as few things as possible go wrong, and assumes that errors occur because of identifiable failures or malfunction of specific components of a system [92]. Safety-II is a more recent initiative that focuses on what is going well in a system or process and analyses the associated everyday working practices with the aim of replicating them more widely [92]. To improve patient safety, it is vital that we learn from the many cases that go well as well as from cases that do not go to plan [93–95].

Debriefs and peer support tools following a death in theatre or critical event

The death of a patient in an operating theatre or a critical event where a patient comes to harm may have a devastating effect on staff involved, even if the event was anticipated [96].

Evidence suggests that a psychological debrief for all staff involved in a critical event, for example critical incident stress debriefing, may encourage staff to relive the trauma and may be harmful [97, 98]. An immediate team ‘check in’, ‘diffusion meeting’ or ‘after action review’ for staff involved may be helpful if short (< 2 min), carried out in a non-judgemental and objective manner and generates a list of staff involved to facilitate supportive follow up [99]. Specific tools may be used by appropriately trained staff to structure this [99].

The use of peer support tools, such as trauma risk management (TRiM) [100], have been used in healthcare [101]. They provide a structure for managing the immediate aftermath of a critical event and can reassure staff that a trauma stress reaction is normal after an abnormal event, and this usually resolves with time [101]. In addition, peer support tools provide a framework for identifying staff who may benefit from professional psychological help after such an event and can signpost them to such services [101].

Education and training

Human factors training

The UK Chartered Institute of Ergonomics and Human Factors white paper [2] and Health Education England’s patient safety syllabus [102] recommend basic human factors awareness training for all NHS staff. In addition, the Ockenden review [103] recommended annual human

factors training for all staff working in a maternity setting. The recommended training includes the role of good design, use of non-technical skills as appropriate for role and an understanding of performance shaping factors [2, 8, 102].

Non-technical skills training

Anaesthetists and their teams can learn and practice non-technical skills during in-theatre training, classroom teaching and by using e-learning programmes. Once the basic concepts are established, simulation training can provide an opportunity for multidisciplinary teams to practise and rehearse non-technical skills and emergency drills/scenarios [73–75, 78]. The use of a dedicated simulation suite allows staff to train away from the distractions of the workplace. Simulation training in an empty operating theatre, recovery bay or clinic room (in-situ simulation) increases realism, enables evaluation of the working environment and working practice design and may prompt improvements [74, 75, 78].

The ANTS system has been used successfully in simulation training centres and provides both a framework to assess non-technical skills and a template for identifying ways to improve them [15, 17, 18, 74]. Similar frameworks have been designed for other staff groups that work with anaesthetists, including: non-technical skills for surgeons (NOTSS) [104]; nurse-anaesthetists' non-technical skills (N-ANTS and NANTS-no) [105, 106]; anaesthetic non-technical skills for anaesthetic practitioners (ANTS-AP) [70]; scrub practitioners' list of intra-operative non-technical skills (SPLINTS) [107]; and the observational teamwork assessment for surgery (OTAS) [108].

Human performance under pressure and cognitive overload

Deterioration in performance in high-pressure situations is well recognised [109] and can be explained by several underlying physiological processes including the 'fight or flight' response [83], neurophysiological response to cognitive overload [72, 79, 83] and 'limbic hijacking' [110].

Activation of the 'fight or flight' response to a minor degree can improve performance, but further activation can lead to prioritisation of gross motor skills over fine motor skills, reducing manual dexterity. In addition, prioritisation of vision over hearing, along with narrowing of the visual field, contribute to poor communication and loss of situation awareness [86].

Working memory has a finite capacity and can typically only hold between four and seven pieces of information at any one time [73, 111]. In a clinical emergency, staff need to process large amounts of information in a short period of

time, with the risk that working memory becomes overwhelmed (cognitive overload) [73, 83, 112] resulting in a fall in performance [73, 83, 112].

Limbic hijacking occurs in high stress situations and results in workers exhibiting impulsive reactions to sensory stimuli [110]; these reactions include the 'startle' or 'surprise' responses and also contribute to a fall in performance [113].

Worker well-being

The importance of worker well-being in maintaining patient safety is increasingly being recognised [2, 4, 97], with the acknowledgement that 'well and healthy staff are safer staff' [2, 4, 97, 114]. The human factors strategies described within the hierarchy of controls model (Fig. 1) improve the well-being of workers as well as the safety of patients [2, 4, 25]. Human factors professionals investigate routinely the interface between workers and their equipment and working environments to reduce the risk of physical injuries. They also use specific tools to investigate the effect of fatigue on worker well-being and safety, and study the impacts of rota design, working hours and rest pauses [2, 114–116]. Such strategies are relevant to current anaesthetic practice [115, 116].

Several high-profile anaesthetic deaths in recent years have shared common themes including: task fixation; loss of situation awareness; steep team hierarchy; poor teamwork and communication; and a fall in personal and team performance in a high-pressure situation [3, 19, 20, 22, 76, 77, 117]. A follow-up study to the 4th National Audit Project (NAP4) [13] examined 12 cases in detail using a human factors investigation tool [118]. Contributing human factors issues were identified in all cases, the most frequent being: loss of situation awareness (failure to anticipate, wrong decision); job factors (task management, staffing, time pressure); and person factors (tiredness, hunger, stress) [118].

While there is no doubt that good non-technical skills and strategies to maintain optimum human performance can contribute to patient safety, these alone cannot compensate for poor working environments, inadequate systems, understaffing and lack of appropriate financial investment. By implementing human factors design principles to avoid error where possible and incorporating barriers and mitigations to prevent errors that do occur from causing harm, the reliance on high levels of personal and team performance currently required in anaesthesia (and healthcare in general) can be reduced and patient safety and worker well-being can be improved.

Acknowledgements

The Difficult Airway Society and the Association of Anaesthetists funded the human factors Working Party meetings and SurveyMonkey subscription which led to the writing of this narrative review. The authors would like to thank M. Bromiley for his advice during the preparation of this narrative review and Dr S. Watson for permission to use a photograph in Fig. 3. NS' research is supported by the National Institute for Health Research (NIHR) Applied Research Collaboration (ARC) South London at King's College Hospital NHS Foundation Trust. NS is a member of King's Improvement Science, which is funded by King's Health Partners (Guy's and St Thomas' NHS Foundation Trust, King's College Hospital NHS Foundation Trust, King's College London and South London and Maudsley NHS Foundation Trust), and the Guy's and St Thomas' Foundation. The views expressed in this publication are those of the author(s) and not necessarily those of the NIHR, KHP, the charities or the Department of Health and Social Care. NS is the director of the London Safety and Training Solutions Ltd, which offers training in patient safety, implementation solutions and human factors to healthcare organisations and the pharmaceutical industry. CF is chair of the Clinical Human Factors Group charity. TC and FK's department has received airway equipment from several manufacturers for research or evaluation either for free or reduced cost. Neither author has benefited financially. TC has lectured, without payment, at an educational meeting of a manufacturer of videolaryngoscopes and attended a meeting, without payment, regarding capnography implementation at another manufacturer of videolaryngoscopes. No other competing interests declared.

References

- International Ergonomics Association. What is ergonomics (HFE)? <https://iea.cc/what-is-ergonomics> (accessed 07/11/2022).
- Chartered Institute of Ergonomics and Human Factors. Human factors in health and social care. 2018. <https://ergonomics.org.uk/resource/human-factors-in-health-and-social-care.html> (accessed 07/11/2022).
- The Health Foundation. Clinical Human Factors Group. https://www.health.org.uk/funding-and-partnerships/programmes/clinical-human-factors-group?gclid=EA1alQobChMl7q-O6eOR-ALVjJftCh0JAzxEAAYASAAEgLRDfD_BwE (accessed 07/11/2022).
- Shorrock S, Williams C. *Human Factors and Ergonomics in Practice: Improving System Performance and Human Wellbeing in the Real World*. New York: CRC Press, 2017.
- Manuele FA. Achieving risk reduction, effectively. *Human Factors Management* 2006; **84**: 184–90.
- Taylor E, Hignett S, Joseph A. The environment of safe care: considering building design as one facet of safety. *Proceedings of the International Symposium of Human Factors and Ergonomics in Healthcare* 2014; **3**: 123–7.
- Ministry of Defence. JSP 912. Human Factors Integration for Defence Systems. Part 2: Guidance. 2021. https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/1061681/JSP912_Part_2.pdf (accessed 07/11/2022).
- Health Education England. Human Factors and Healthcare: Evidencing the impact of Human Factors training to support improvements in patient safety and to contribute to cultural change. 2019. <https://www.hee.nhs.uk/sites/default/files/documents/Health%20Education%20England%20and%20CIEHF%20-%20Human%20Factors%20and%20Healthcare%20Report.pdf> (accessed 07/11/2022).
- Helmreich RL, Merritt AC, Wilhelm JA. The evolution of crew resource management training in commercial aviation. *International Journal of Aviation* 1999; **9**: 19–32.
- Centers for Disease Control and Prevention. Hierarchy of controls. 2022. <https://www.cdc.gov/niosh/topics/hierarchy/> (accessed 07/11/2022).
- Libertati EG, Perrall MF, Dixon WM. Learning from high-risk industries may not be straightforward: a qualitative study of the hierarchy of risk controls approach in healthcare. *International Journal for Quality in Health Care* 2018; **30**: 39–43.
- Reason J. Human error: models and management. *British Medical Journal* 2000; **320**: 768–70.
- Royal College of Anaesthetists/Difficult Airway Society. Major complications of airway management in the United Kingdom. 4th National Audit Project. 2011. <https://www.nationalauditprojects.org.uk/downloads/NAP4%20Full%20Report.pdf> (accessed 07/11/2022).
- Flin R, O'Connor P, Crichton M. *Safety at the Sharp End: A Guide to Non-Technical Skills*. Farnham, UK: Ashgate, 2008.
- Flin R, Patey R, Glavin R, Maran N. Anaesthetists' non-technical skills. *British Journal of Anaesthesia* 2010; **105**: 38–44.
- Howard S, Gaba D, Fish K, Yang G, Sarnquist FH. Anesthesia crisis resource management training: teaching anesthesiologists to handle critical incidents. *Aviation, Space, and Environmental Medicine* 1992; **63**: 763–70.
- Fletcher G, McGeorge P, Flin R, et al. The role of non-technical skills in anaesthesia: a review of current literature. *British Journal of Anaesthesia* 2002; **88**: 418–29.
- Fletcher G, Flin R, McGeorge P, Glavin R, Maran N, Patey R. Anaesthetists' non-technical skills (ANTS): evaluation of a behavioural marker system. *British Journal of Anaesthesia* 2003; **90**: 580–8.
- Bromiley M. Have you ever made a mistake? *Royal College of Anaesthetists Bulletin* 2008; **48**: 2442–5. <https://studylib.net/doc/18282697/have-you-ever-made-a-mistake%3F> (accessed 07/11/2022).
- New Statesman. How mistakes can save lives: one man's mission to revolutionise the NHS. 2014. <https://www.newstatesman.com/long-reads/2014/06/how-mistakes-can-save-lives> (accessed 07/11/2022).
- Dul J, Bruder R, Buckle P, et al. A strategy for human factors/ergonomics: developing the discipline and profession. *Ergonomics* 2012; **55**: 377–95.
- Milton Keynes Coroner's Court. Inquest into the death of Glenda May Logsdail. 2021. https://www.judiciary.uk/wp-content/uploads/2021/09/Glenda-Logsdail-Prevention-of-future-deaths-report-2021-0295_Published.pdf (accessed 03/06/2022).
- Royal College of Anaesthetists. Patient safety: unrecognised oesophageal intubation. <https://www.rcoa.ac.uk/safety-standards-quality/patient-safety/prevention-future-deaths> (accessed 07/11/2022).
- Griffiths R, Wells A, Logsdail R. Why does oesophageal intubation still go unrecognised? *Anaesthesia* 2022; **77**: 723–4.

25. Sujan M, Pickup L, Vosper H, et al. Making human factors and ergonomics work in health and social care: chapter 1. <https://ergonomics.org.uk/resource/hf-in-health-and-social-care-ebook-chapter-1.html> (accessed 7/11/22).
26. Hansel J, Rogers AM, Lewis SR, Cook TM, Smith AF. Videolaryngoscopy versus direct laryngoscopy for adults undergoing tracheal intubation. *Cochrane Database of Systematic Reviews* 2022; **4**: CD011136.
27. Lewis SR, Butler AR, Parker J. Videolaryngoscopy versus direct laryngoscopy for adult patients requiring tracheal intubation: a Cochrane systematic review. *British Journal of Anaesthesia* 2017; **119**: 369–83.
28. Grundgeiger T, Roewer N, Grundgeiger J, Hurtienne J, Happel O. Body posture during simulated tracheal intubation: Glidescope (®) videolaryngoscopy vs Macintosh direct laryngoscopy for novices and experts. *Anaesthesia* 2015; **70**: 1375–81.
29. Hall D, Steel A, Heij R, Eley A, Young P. Videolaryngoscopy increases 'mouth-to-mouth' distance compared with direct laryngoscopy. *Anaesthesia* 2020; **75**: 822–3.
30. Sanfilippo F, Tigano S, La Rosa V, et al. Tracheal intubation while wearing personal protective equipment in simulation studies: a systematic review and meta-analysis with trial-sequential analysis. *Brazilian Journal of Anesthesiology* 2022; **72**: 291–301.
31. International Electrotechnical Commission. IEC 62366-1:2015 - Part 1: Application of usability engineering to medical devices. International Electrotechnical Commission. 2015. <https://www.iso.org/standard/63179.html> (accessed 07/11/2022).
32. International Electrotechnical Commission. IEC/TR 62366-2 - Part 2: Guidance on the application of usability engineering to medical devices. International Electrotechnical Commission. 2016. <https://www.iso.org/standard/69126.html> (accessed 07/11/2022).
33. Medicines and Healthcare Products Regulatory Agency. Guidance on applying human factors to medical devices. 2017 (updated 2021). <https://www.gov.uk/government/publications/guidance-on-applying-human-factors-to-medical-devices> (accessed 07/11/2022).
34. Newton R, Mytton OT, Aggarwal R. Making existing technology safer in healthcare. *British Medical Journal Quality and Safety* 2010; **19**: i15–i24.
35. American Society of Anesthesiologists. Statement on Labeling of Pharmaceuticals for Use in Anesthesiology. 2004 (amended 2020). <https://www.asahq.org/standards-and-guidelines/statement-on-labeling-of-pharmaceuticals-for-use-in-anesthesiology> (accessed 07/11/2022).
36. International Organization of Standardization. ISO 26825:2008 Anaesthetic and respiratory equipment — User-applied labels for syringes containing drugs used during anaesthesia — Colours, design and performance. 2008. <https://www.iso.org/standard/76678.html> (accessed 7/11/22).
37. Medicines and Healthcare Products Regulatory Agency. Best practice in the labelling and packaging of medicines. 2014 (updated 2020). <https://www.gov.uk/government/publications/best-practice-in-the-labelling-and-packaging-of-medicines> (accessed 07/11/2022).
38. Keers R, Williams S, Cooke J, et al. Prevalence and nature of medication administration errors in health care settings: a systematic review of direct observational evidence. *Annals of Pharmacotherapy* 2013; **47**: 237–56.
39. Oglesby FC, Ray AG, Shurlock T, Mitra T, Cook TM. Litigation related to anaesthesia: analysis of claims against the NHS in England 2008–2018 and comparison against previous claim patterns. *Anaesthesia* 2022; **77**: 527–37.
40. Cranshaw J, Gupta KJ, Cook TM. Litigation related to drug errors in anaesthesia: an analysis of claims against the NHS in England 1995–2007. *Anaesthesia* 2009; **64**: 1317–23.
41. Almgairbi L, Sharp R, Griffiths R, et al. An observational feasibility study of a new anaesthesia drug storage tray. *Anaesthesia* 2018; **73**: 356–64.
42. Hellier E, Edworthy J, Derbyshire N, Costello A. Considering the impact of medicine label design characteristics on patient safety. *Ergonomics* 2006; **49**: 617–30.
43. Bailey CR, Radhakrishna S, Asanati K, et al. Ergonomics in the anaesthetic workplace: guideline from the Association of Anaesthetists. *Anaesthesia* 2019; **76**: 1635–47.
44. Jurewicz KA, Neyens DM, Catchpole K, Joseph A, Reeves ST, Abernathy JH III. Observational study of anaesthesia workflow to evaluate physical workspace design and layout. *British Journal of Anaesthesia* 2021; **126**: 633–41.
45. Schnittker R, Marshall S, Horberry T, Young KL. Human factors enablers and barriers for successful airway management – an in-depth interview study. *Anaesthesia* 2018; **73**: 980–9.
46. Walker ST, Brett SJ, McKay A, Aggarwal R, Vincent C. 'The Resus Station': the use of clinical simulations in a randomised crossover study to evaluate a novel resuscitation trolley. *Resuscitation* 2012; **83**: 1374–80.
47. Frerk C, Mitchell VS, McNarry AF, et al. Difficult airway society 2015 guidelines for management of unanticipated difficult intubation in adults. *British Journal of Anaesthesia* 2015; **115**: 827–48.
48. Chan J, Shojania KG, Easty AC, Etchells EE. Does user-centred design affect the efficiency, usability and safety of CPOE order sets? *Journal of the American Medical Informatics Association* 2011; **18**: 276–81.
49. Federal Register. Health Information Technology: Standards, Implementation Specifications, and Certification Criteria for Electronic Health Record Technology, 2014 Edition; Revisions to the Permanent Certification Program for Health Information Technology. 2012. <https://federalregister.gov/a/2012-20982> (accessed 07/11/2022).
50. Sasangohar F, Peres SC, Williams J, et al. Investigating written procedures in process safety: qualitative data analysis of interviews from high-risk facilities. *Process Safety and Environmental Protection* 2018; **113**: 30–9.
51. Vincent C, Amalberti R. *Safer Healthcare: Strategies for the Real World*. London: Springer, 2016.
52. Reiley S, Grasha AF, Schafer J. Workload, error detection, and experienced stress in a simulated pharmacy verification task. *Perceptual and Motor Skills* 2002; **95**: 27–46.
53. Arora S, Sevdalis N, Nestel D, Woloshynowych M, Darzi A, Kneebone R. The impact of stress on surgical performance: a systematic review of the literature. *Surgery* 2010; **147**: 318–30.
54. Arora S, Sevdalis N, Aggarwal R, Sirimanna P, Darzi A, Kneebone R. Stress impairs psychomotor performance in novice laparoscopic surgeons. *Surgical Endoscopy* 2010; **24**: 2588–93.
55. Arora S, Hull L, Sevdalis N, et al. Factors compromising safety in surgery: stressful events in the operating room. *American Journal of Surgery* 2010; **199**: 60–5.
56. McElhatton J, Drew C. Time pressure as a causal factor in aviation safety incidents - The 'hurry-up' syndrome. International Symposium on Aviation Psychology 1993. https://asrs.arc.nasa.gov/publications/directline/dl5_hurry.htm (accessed 07/11/2022).
57. Leviatan I, Oberman B, Zimlichman E, Stein GY. Associations of physicians' prescribing experience, work hours, and workload with prescription errors. *Journal of the American Medical Informatics Association* 2021; **28**: 1074–80.

58. Pandit JJ, The NHS. Improvement report on operating theatres: really 'getting it right first time'? *Anaesthesia* 2019; **74**: 839–44.
59. Marshall S. The use of cognitive aids during emergencies in anaesthesia: a review of the literature. *Anesthesia and Analgesia* 2013; **117**: 1162–71.
60. Reason J. Cognitive aids in process environments: prostheses or tools? *International Journal of Man-Machine Studies* 1987; **27**: 463–70.
61. Association of Anaesthetists. Quick Reference Handbook. Updated 2022. <https://www.anesthesiologists.org/qrh> (accessed 07/11/2022).
62. Chrimes N. The vortex: a universal 'high-acuity implementation tool' for emergency airway management. *British Journal of Anaesthesia* 2016; **117**: i20–i27.
63. Haynes AB, Weiser TG, Berry WR, et al. A surgical safety checklist to reduce morbidity and mortality in a global population. *New England Journal of Medicine* 2009; **360**: 491–9.
64. Treadwell JR, Lucas S, Tsou AY. Surgical checklists: a systematic review of impacts and implementation. *British Medical Journal Quality and Safety* 2014; **23**: 299–318.
65. Haugen AS, Sevdalis N, Søfteland E. Impact of the World Health Organization surgical safety checklist on patient safety. *Anesthesiology* 2019; **131**: 420–5.
66. Cook TM, El-Boghdady K, McGuire B, et al. Consensus guidelines for managing the airway in patients with COVID-19: guidelines from the difficult airway society, the Association of Anaesthetists the Intensive Care Society, the Faculty of Intensive Care Medicine and the Royal College of Anaesthetists. *Anaesthesia* 2020; **75**: 785–99.
67. Higgs A, McGrath BA, Goddard C, et al. Guidelines for the management of tracheal intubation in critically ill adults. *British Journal of Anaesthesia* 2018; **120**: 323–52.
68. Royal College of Anaesthetists. Curriculum for a CCT in Anaesthetics. 2010. <https://www.rcoa.ac.uk/sites/default/files/documents/2019-08/TRG-CU-CCT-ANAES2010.pdf> (accessed 07/11/2022).
69. Fadden S, Mercer S. Followership in complex trauma. *Trauma* 2019; **21**: 6–13.
70. Rutherford JS, Flin R, Irwin A, McFadyen AK. Evaluation of the prototype Anaesthetic Non-technical Skills for Anaesthetic Practitioners (ANTS-AP) system: a behavioural rating system to assess the non-technical skills used by staff assisting the anaesthetist. *Anaesthesia* 2015; **70**: 907–14.
71. Anaesthetists' non-technical skills (ANTS) system handbook v1.0. 2012. <https://research.abdn.ac.uk/wp-content/uploads/sites/14/2019/03/ANTS-Handbook-2012-1.pdf> (accessed 7/11/2022).
72. Schulz C, Endsley M, Kochs E, et al. Situation awareness in anesthesia: concept and research. *Anesthesiology* 2013; **118**: 729–42.
73. Wickens CD. Multiple resources and performance prediction. *Theoretical Issues in Ergonomics Science* 2002; **3**: 159–77.
74. Manser T, Harrison TK, Gaba DM, Howard SK. Coordination patterns related to high clinical performance in a simulated anesthetic crisis. *Anesthesia and Analgesia* 2009; **108**: 1606–15.
75. Schmutz J, Meier L, Manser T. How effective is teamwork really? The relationship between teamwork and performance in healthcare teams: a systematic review and meta-analysis. *British Medical Journal Open* 2019; **9**: e028280.
76. Courts and Tribunals Judiciary. Investigation into the death of Sharon Grierson. 2018. https://www.judiciary.uk/wp-content/uploads/2018/06/Sharon-Grierson-2018-0034_Redacted.pdf (accessed 07/11/2022).
77. Courts and Tribunals Judiciary. Investigation into the death of Peter Saint. 2017. https://www.judiciary.uk/wp-content/uploads/2018/02/Peter-Saint-2017-0404_Redacted.pdf (accessed 07/11/2022).
78. Boyd M, Cumin D, Lombard Boyd M, et al. Read-back improves information transfer in simulated clinical crises. *British Medical Journal Quality and Safety* 2014; **23**: 989–93.
79. Härgestam M, Hultin M, Brulin C, Jacobsson M. Trauma team leaders' non-verbal communication: video registration during trauma team training. *Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine* 2016; **24**: 1–10.
80. Zala-Mező E, Wacker J, Künzle B, et al. The influence of standardisation and task load on team coordination patterns during anaesthesia inductions. *British Medical Journal Quality and Safety* 2009; **18**: 127–30.
81. Sevdalis N, Undre S, McDermott J, Giddie J, Diner L, Smith G. Impact of intraoperative distractions on patient safety: a prospective descriptive study using validated instruments. *World Journal of Surgery* 2014; **38**: 751–8.
82. Van Harten A, Gooszen HG, Koksmo JJ, et al. An observational study of distractions in the operating theatre. *Anaesthesia* 2021; **76**: 346–56.
83. Hearns S. *Peak Performance Under Pressure*. Somerset, UK: Class Professional Publishing, 2019.
84. Patini N, Arzola C, Malavade A, Varmani S, Krimus L, Friedman Z. Challenging authority and speaking up in the operating room environment: a narrative synthesis. *British Journal of Anaesthesia* 2019; **122**: 223–44.
85. Weller JM, Long JA. Creating a climate for speaking up. *British Journal of Anaesthesia* 2019; **122**: 710–3.
86. Eppich W. "Speaking up" for patient safety in the pediatric emergency department. *Clinical Pediatric Emergency Medicine* 2015; **16**: 83–9.
87. Fioratou E, Flin R, Glavin R. No simple fix for fixation errors: cognitive processes and their clinical applications. *Anaesthesia* 2010; **65**: 61–9.
88. NHS Improvement. Surgical Never Events Learning from 38 cases occurring in English hospitals between April 2016 and March 2017. 2018. https://spiral.imperial.ac.uk/bitstream/10044/1/72410/2/Learning_from_surgical_Never_Events_FINAL.pdf (accessed 07/11/2022).
89. NHS England. Patient Safety Incident Response Framework. <https://www.england.nhs.uk/patient-safety/incident-response-framework/> (accessed 07/11/2022).
90. Holden R, Carayon P. SEIPS 101 and seven simple SEIPS tools. *British Medical Journal Quality and Safety* 2021; **30**: 901–10.
91. Lawton R, McEachan RRC, Giles SJ, et al. Development of an evidence-based framework of factors contributing to patient safety incidents in hospital settings: a systematic review. *British Medical Journal Quality and Safety* 2012; **21**: 369–80.
92. Hollnagel E. *Safety—I and Safety—II: The Past and Future of Safety Management*. Boca Raton, FL: CRC Press, 2018.
93. Mitchell EL, Lee DY, Arora S, et al. Improving the quality of the surgical morbidity and mortality conference: a prospective intervention study. *Academy Medicine* 2013; **88**: 824–30.
94. Kelly N, Blake S, Plunkett A. Learning from excellence in healthcare: a new approach to incident reporting. *Archives of Disease in Childhood* 2016; **101**: 788–91.
95. Health Improvement Scotland. Scottish Mortality and Morbidity Programme. https://www.healthcareimprovementscotland.org/our_work/patient_safety/scottish_mortality_morbidity.aspx (accessed 07/11/2022).
96. Royal College of Anaesthetists. Chapter 5: Guidelines for the Provision of Emergency Anaesthesia Services. 2022. <https://www.rcoa.ac.uk/gpas/chapter-5> (accessed 07/11/2022).
97. Brooks SK, Rubin GJ, Greenberg N. Traumatic stress within disaster-exposed occupations: overview of the literature and suggestions for the management of traumatic stress in the workplace. *British Medical Bulletin* 2019; **129**: 25–34.

98. National Institute of Health and Care Excellence. Post-traumatic stress disorder. [NG116]. 2018. <https://www.nice.org.uk/guidance/ng116> (accessed 07/11/2022).
99. Johnson TJ, Millinchamp FJ, Kelly FE. Use of a team immediate debrief tool to improve staff well-being after potentially traumatic events. *Anaesthesia* 2019; **76**: 1001–2.
100. Greenberg N, Langston V, Jones N. Trauma risk management (TRIM) in the UK armed forces. *Journal of the Royal Army Medical Corps* 2008; **154**: 124–7.
101. Kelly FE, Osborn M, Stacey MS. Improving resilience in anaesthesia and intensive care medicine – learning lessons from the military. *Anaesthesia* 2020; **75**: 720–3.
102. Health Education England. Patient safety. <https://www.hee.nhs.uk/our-work/patient-safety> (accessed 07/11/2022).
103. Department of Health and Social Care. Final report of the Ockenden review. 2022. <https://www.gov.uk/government/publications/final-report-of-the-ockenden-review> (accessed 07/11/2022).
104. Yule S, Flin R, Maran N, Rowley D, Youngson G, Paterson-Brown S. Surgeons' non-technical skills in the operating room: reliability testing of the NOTSS behaviour rating system. *World Journal of Surgery* 2008; **32**: 548–56.
105. Lyk-Jensen HT, Jepsen RMHG, Spanager L, et al. Assessing nurse Anaesthetists' non-technical skills in the operating room. *Acta Anaesthesiologica Scandinavica* 2014; **58**: 794–801.
106. Flynn F, Valeberg B, Tønnessen S, et al. Psychometric testing of a structured assessment instrument for non-technical skills (NANTS-no) for use in clinical supervision of student nurse anesthetists. *Journal of Nursing Measurement* 2020; **291**: E59–E77.
107. Mitchell L, Flin R, Yule S, Mitchell J, Coutts K, Youngson G. Evaluation of the scrub practitioners' list of intraoperative non-technical skills (SPLINTS) system. *International Journal of Nursing Studies* 2012; **49**: 201–11.
108. Undre S, Sevdalis N, Healey AN, Darzi A, Vincent CA. Observational teamwork assessment for surgery (OTAS): refinement and application in urological surgery. *World Journal of Surgery* 2007; **31**: 1373–81.
109. Yerkes RM, Dodson JD. The relation of strength of stimulus to rapidity of habit-formation. *Journal of Comparative Neurology and Psychology* 1908; **18**: 459–82.
110. Goleman D. *Emotional Intelligence: Why It Can Matter More Than IQ*. London: Bloomsbury Publishing, 1995.
111. Cowan N. The magical mystery four: how is working memory capacity limited, and why? *Current Directions in Psychological Science* 2010; **19**: 51–7.
112. Byrne AJ, Oliver M, Bodger O. Novel method of measuring the mental workload of anaesthetists during clinical practice. *British Journal of Anaesthesia* 2010; **6**: 767–71.
113. Rivera J, Talone AB, Boesser CT, Jentsch F, Yeh M. Startle and surprise on the flight desk: similarities, differences and prevalence. *Proceedings of the Human Factors and Ergonomics Society Annual Meeting* 2014; **58**: 1047–51.
114. General Medical Council. Caring for doctors, caring for patients. 2019. https://www.gmc-uk.org/-/media/documents/caring-for-doctors-caring-for-patients_pdf-80706341.pdf (accessed 07/11/2022).
115. Association of Anaesthetists. Help Fight Fatigue. <https://anaesthetists.org/Home/Wellbeing-support/Fatigue/Help-Fight-Fatigue> (accessed 07/11/2022).
116. Association of Anaesthetists. Wellbeing and support. <https://anaesthetists.org/Home/Wellbeing-support> (accessed 07/11/2022).
117. Scottish Courts and Tribunals. Inquiry under the fatal accidents and inquiries (Scotland) Act 1976 into the sudden death of Gordon Ewing. 2010. <https://www.scotcourts.gov.uk/search-judgments/judgment?id=328e86a6-8980-69d2-b500-ff0000d74aa7> (accessed 07/11/2022).
118. Flin R, Fioratou C, Frerk C, et al. Human factors in the development of complications of airway management: preliminary evaluation of an interview tool. *Anaesthesia* 2013; **68**: 817–25.